

MULTISITE OPTICAL IMAGING OF ARTIFICIAL IONOSPHERIC PLASMAS (POSTPRINT)

Todd R. Pedersen, et al.

09 November 2011

Interim Report

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.



**AIR FORCE RESEARCH LABORATORY
Space Vehicles Directorate
3550 Aberdeen Ave SE
AIR FORCE MATERIEL COMMAND
KIRTLAND AIR FORCE BASE, NM 87117-5776**

DTIC COPY

NOTICE AND SIGNATURE PAGE

Using Government drawings, specifications, or other data included in this document for any purpose other than Government procurement does not in any way obligate the U.S. Government. The fact that the Government formulated or supplied the drawings, specifications, or other data does not license the holder or any other person or corporation; or convey any rights or permission to manufacture, use, or sell any patented invention that may relate to them.

This report was cleared for public release by the 66 ABW Public Affairs Office and is available to the general public, including foreign nationals. Copies may be obtained from the Defense Technical Information Center (DTIC) (<http://www.dtic.mil>).

AFRL-RV-PS-TR-2013-0113 HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION IN ACCORDANCE WITH ASSIGNED DISTRIBUTION STATEMENT.

//SIGNED//

Daniel G. Miller
Program Manager, AFRL/RVBXI

//SIGNED//

Edward J. Masterson, Colonel, USAF
Chief, Battlespace Environment Division

This report is published in the interest of scientific and technical information exchange, and its publication does not constitute the Government's approval or disapproval of its ideas or findings.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 09-11-2011		2. REPORT TYPE Interim Report		3. DATES COVERED (From - To) Nov 2010 to Nov 2011	
4. TITLE AND SUBTITLE Multisite Optical Imaging of Artificial Ionospheric Plasmas (Postprint)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 61102F	
6. AUTHOR(S) Todd R. Pedersen, Jeffrey M. Holmes, Bjorn Gustavsson, and Travis J. Mills				5d. PROJECT NUMBER 3001	
				5e. TASK NUMBER PPM00005508	
				5f. WORK UNIT NUMBER EF004372	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory Space Vehicles Directorate 3550 Aberdeen Avenue SE Kirtland AFB, NM 87117-5776				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-RV-PS-TR-2013-0113	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/RVBXI	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. Approved for public release; distribution is unlimited. (66ABW-2010-1304 dtd Nov 2010)					
13. SUPPLEMENTARY NOTES Government Purpose Rights IEEE Transactions on Plasma Science, 9 November 2011					
14. ABSTRACT Artificial ionospheric plasmas are formed on the bottom side of the natural ionospheric F region during high-power high-frequency (HF) heating experiments and descend to altitudes as low as 140 km before disappearing. Optical emissions produced during these events often exhibit bull's-eye structures, where the artificial plasma is thought to form a central spot that diverts or blocks HF waves to form an empty ring of emissions from the natural ionosphere at higher altitudes. We present multisite image data showing that, in some cases, both the spot and ring represent distinct artificial plasma layers.					
15. SUBJECT TERMS Ionosphere, optical imaging, plasma density, radio propagation					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unlimited	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON Daniel G. Miller
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

This page is intentionally left blank.

Multisite Optical Imaging of Artificial Ionospheric Plasmas

Todd R. Pedersen, Jeffrey M. Holmes, Bjorn Gustavsson, and Travis J. Mills

Abstract—Artificial ionospheric plasmas are formed on the bottom side of the natural ionospheric F region during high-power high-frequency (HF) heating experiments and descend to altitudes as low as 140 km before disappearing. Optical emissions produced during these events often exhibit bull's-eye structures, where the artificial plasma is thought to form a central spot that diverts or blocks HF waves to form an empty ring of emissions from the natural ionosphere at higher altitudes. We present multisite image data showing that, in some cases, both the spot and ring represent distinct artificial plasma layers.

Index Terms—Ionosphere, optical imaging, plasma density, radio propagation.

ARTIFICIAL IONOSPHERIC plasmas are localized regions of enhanced plasma density observed descending from the background natural ionosphere within the beam of a high-power high-frequency (HF) transmitter, particularly when the transmitter frequency is near the second harmonic of the electron gyrofrequency ($2f_{ce}$). Recently discovered at the High-Frequency Active Auroral Research Program (HAARP) facility in Gakona, Alaska (62.4° N 145° W) after the transmitter reached full 3.6-MW power, these plasmas appear in ionosonde measurements as distinct layers of echoes below those from the natural ionosphere, in regions of otherwise much lower plasma density [1]. Although the plasma production mechanism remains uncertain, a critical aspect of the artificial plasmas is that they are dense enough to interact directly with the HF waves and draw energy from the transmitter beam until eventually losing critical density and disappearing at altitudes near 150 km [1], [2]. Absorption of power within the plasma results in enhanced yet generally subvisual optical emissions excited by the impact of thermal and accelerated electrons [3]. Imaging of these emissions from multiple ground sites played a key role in the discovery of the plasmas and in the realization that they were self-sustaining layers dependent on the transmitter. In particular, in [1], optical emissions formed

a bull's-eye pattern when seen from below, but remote images showed the artificial plasma in the center of the bull's-eye to be ~50 km below the ring, which resulted from excitation of the natural ionosphere. In this paper, we present multisite images of an artificial ionospheric plasma which shows both the spot and ring to be distinct artificial layers.

The experiment was carried out on November 19, 2009, between 02:26 UT and 02:43:50 UT. Optical images were acquired at the HAARP site at 557.7 nm (O ¹S, excited by electrons > 4.19 eV) with a 1" Princeton Instruments Versarray 1340 × 1300 pixel camera equipped with all-sky [180° field of view (FOV)] telecentric optics built by Keo Consultants and a 2-nm narrow-band filter. The detector was binned to 335 × 325 pixel resolution to enhance signal to noise and integrated for 5 s at a temperature of −40 °C. A second system located 160 km north of the HAARP near Delta Junction used an Apogee Alta E47 1/2" 1024 × 1024 pixel CCD detector binned to 256 × 256 pixels and integrated for 13 s at −40 °C. This system viewed a field-broadening mirror through a 50-mm f/1.2 Nikon lens and a 10-nm filter centered at 560 nm, providing an effective FOV of ~70°, which was centered at 45° elevation 180° azimuth. The HAARP HF transmitter was pointed vertically and operated at 3.6 MW (~440 MW effective radiated power) in a frequency sweep mode from 2.85 to 2.95 MHz designed to match $2f_{ce}$ in the artificial plasma as it descended to altitudes of stronger magnetic field and higher gyrofrequency. The transmitter parameters and plasma density contours for this period are available in [2].

Fig. 1 shows a schematic of the observation setup and image data, shown in false color approximating the actual wavelength, from the remote system looking obliquely (upper left, acquired at 02:36:15 UT) and the HAARP imager looking up (top center, acquired at 02:36:07 UT). Note that the actual FOV in the image data is much broader than the notional lines of sight in the diagram. The up-looking image from the HAARP shows a partial but well-defined spot and a concentric ring separated by a pronounced dark gap. The oblique image of the same structure also shows a clear gap, but the gap is much wider at the top of the image and disappears near the bottom. This is consistent with a geometry similar to an inverted top hat, where the central spot forms a lower layer corresponding to the top of the hat, while the ring forms another layer at higher altitude corresponding to the brim of the hat. Triangulation of multiple points within both the ring and the spot over ~15 min of the experiment shows the spot to be ~15 km below the ring, a separation which was maintained from the initial formation near 200-km altitude to the final decay at 145 km as the structure descended steadily at ~75 m/s. At the time of these

Manuscript received November 19, 2010; revised April 14, 2011; accepted April 24, 2011. This work was supported in part by the Air Force Office of Scientific Research and in part by the Space Vehicles Directorate Innovation Fund.

T. R. Pedersen, J. M. Holmes, and T. J. Mills are with the Air Force Research Laboratory, Space Vehicles Directorate, Hanscom AFB, MA 01731 USA (e-mail: AFRL.RVB.pa@hanscom.af.mil).

B. Gustavsson is with the University of Southampton, SO171BJ Southampton, U.K. (e-mail: bjorn@irf.se).

Digital Object Identifier 10.1109/TPS.2011.2151209




Fig. 1. Observation geometry and image data from two low-light imaging systems capturing 557.7-nm emissions from a bull's-eye-shaped artificial ionospheric plasma over the HAARP facility. A reconstruction based on the image data shows the central spot and the ring to form two distinct artificial layers separated in altitude by ~ 15 km, which matches closely the multiple layers seen in ionosonde echoes (lower right).

images, the spot altitude was ~ 160 km, and the ring was at ~ 175 km. Fitting of circles to the spot, gap, and ring in the up-looking image gives radii of $\sim 6.4^\circ$ for the edge of the spot, $\sim 8^\circ$ for the inner edge of the ring, and $\sim 13^\circ$ for the overall distribution. These compare to the 10° N-S and 8° E-W half-power half widths for the transmitter beam, shown in blue in the diagram. At the triangulated 160–175-km altitude for the images shown, the central spot is ~ 40 km in diameter, while the inner edge of the ring is ~ 50 km in diameter, and the outer edge of the entire pattern is ~ 80 km in diameter.

The fact that the dark gap remains distinct even though it is less than 2° or ~ 5 km wide indicates that the optical emissions originate in relatively thin layers and do not have large extent along the magnetic field. The inverted top-hat geometry reconstructed from these optical images sheds new light on the ionosonde data (lower right), which showed two artificial layers ~ 15 – 20 km apart in altitude during this period.

In conclusion, multisite optical imaging has demonstrated that artificial ionospheric plasmas can take the form of two distinct layers in a 3-D inverted top-hat configuration. Accurate determination of the properties of such plasmas is essential for the development of practical applications such as enhanced over-the-horizon radar propagation.

REFERENCES

- [1] T. Pedersen, B. Gustavsson, E. Mishin, E. Kendall, T. Mills, H. C. Carlson, and A. L. Snyder, "Creation of artificial ionospheric layers using high-power HF waves," *Geophys. Res. Lett.*, vol. 37, p. L02106, 2010.
- [2] T. Pedersen, M. McCarrick, B. Reinisch, B. Watkins, R. Hamel, and V. Paznukhov, "Production of artificial ionospheric layers by frequency sweeping near the 2nd gyroharmonic," *Ann. Geophys.*, vol. 29, pp. 47–51, 2011.
- [3] P. A. Bernhardt, C. A. Selcher, C. L. Siefring, and E. Gerken, "Imaging of ionospheric density structures and plasma drifts using artificial illumination by high-power radio waves," *IEEE Trans. Plasma Sci.*, vol. 33, no. 2, pp. 504–505, Apr. 2005.

DISTRIBUTION LIST

DTIC/OCF	
8725 John J. Kingman Rd, Suite 0944	
Ft Belvoir, VA 22060-6218	1 cy
AFRL/RVIL	
Kirtland AFB, NM 87117-5776	2 cys
Official Record Copy	
AFRL/RVBXI/Daniel G. Miller	1 cy

This page is intentionally left blank.